

Annapolis Pediatrics Physical Guide

Age	Birth	7-10 days	1 month	2 months	4 months	6 months	9 months	12 months ^c	15 months	18 months	2 years	30 month	3 years	4 years	5 years	6 years	7-8 years	9 years	10 years	11-12 years	13-15 years	16 years	17 years
Labwork	Newborn Screen #1	Newborn Screen #2						Lead/ Hgb			Lead/ Hgb							Cholesterol #1 ^e				GC/ Chlamydia	*females only ^f
																							Cholesterol #2
Physicals	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v
Screening			Edinburgh	Edinburgh	Edinburgh	Edinburgh	ASQ			ASQ & MCHAT	ASQ & MCHAT	ASQ	ASQ	Hearing/ Vision				<-----PHQ9----->					
Vaccines																							
Hepatitis B	#1		#2 ^a				#3																
Rotavirus				#1	#2	#3																	
DTaP				#1	#2	#3																	
Polio (IPV)				#1	#2	#3																	
HIB				#1	#2	#3																	
Pneumococcal				#1	#2	#3		#4															
Influenza						#1 ^b	b	b	b	b	b	b	b	b	b	b	b	b	b	b	b	b	b
MMR								#1 ^c															
Varicella								#1 ^c															
Hepatitis A									#1		#2												
Meningococcal																					#1		#2
HPV																							#1 ^e
Tdap																							#1

Revised 1/12/21

- a - Hepatitis B #2 must be given at least 28 days after Hepatitis B #1, which is typically given at birth.
 - b - Must be at least 6 months of age to receive the influenza vaccine. If under 9 years of age and receiving first dose EVER, patient needs a booster dose within the same season 28 days later.
Influenza vaccine is recommended yearly.
 - c - Must have reached 4th birthday in order to receive vaccines. If not given at 4 years, catchup by 6 years
 - d - May have additional dose of Inactivated Poliovirus at this age.
 - e - Gardasil is approved for males&females at age 9 but typically given at 11 year visit. If started before 15 years of age, Dose #1 and #2 administered 6-12 months apart.
If first dose of HPV is administered after patient turns 15 years old, 3 doses are needed (2nd dose 2 months after first, 3rd dose 4 months after second AND at least 6 months after first).
 - f- starting at age 16, a urine is collected from all female patients for chlamydia and gonorrhea screening for sexually active pts
 - g- if cholesterol not performed at 9 year visit, catch-up screening done between 9&11 years
- Edinburgh-** Depression screening for postpartum women.
- ASQ-** Developmental screening, specifically assessing skills in these areas: communication, gross motor, fine motor, problem solving, personal-social.
- MCHAT-** Developmental screening used to identify children who may benefit from a more thorough developmental and autism evaluation.
- PHQ9-** Multipurpose mental health screening used to diagnose, measure, and monitor the severity of depression in adolescent patients.

